

Historic, archived document

Do not assume content reflects current
scientific knowledge, policies, or practices.

Ag. 84M
Cop. 2

Supplement I:
Annotated Bibliography of the Horn Fly,
***Haematobia irritans irritans* (L.)**
Including References on the Buffalo Fly,
***H. irritans exigua* (de Meijere), and Other**
Species Belonging to the Genus *Haematobia*


Miscellaneous Publication No. 1278

PROCUREMENT SECTION
CURRENT SERIAL RECORDS

APR 20 '77

U.S. DEPT. OF AGRICULTURE
NATL. AGRIC. LIBRARY
RECEIVED

Agricultural Research Service
UNITED STATES DEPARTMENT OF AGRICULTURE



This publication reports research involving pesticides. It does not contain recommendations for their use, nor does it imply that the uses discussed here have been registered. All uses of pesticides must be registered by appropriate State and/or Federal agencies before they can be recommended.

CAUTION: Pesticides can be injurious to humans, domestic animals, beneficial insects, desirable plants, and fish or other wildlife—if they are not handled or applied properly. Use all pesticides selectively and carefully. Follow recommended practices for the disposal of surplus pesticides and pesticide containers.



Use Pesticides Safely
FOLLOW THE LABEL

U.S. DEPARTMENT OF AGRICULTURE

SPECIES CITED IN REFERENCES

- Haematobia* Lepeletier & Serville (1828)
Lyperosia Rondani (1856)
Priophora Robineau-Desvoidy (1863)
Glossinella Grunberg (1906)
Haphospatha Enderlein (1924)
Siphona auct. (nec Meigen)
 irritans irritans (Linnaeus, 1758)
 pungens (Fabricius, 1794)
 serrata Robineau-Desvoidy (1830)
 tibialis Robineau-Desvoidy (1830)
 cornicola Williston (1889)
 meridionalis (Bezzi, 1911)
 ruffifrons (Bezzi, 1911)
 weissi (Bezzi, 1911)
 irritans exigua de Meijere (1903)
 flavohirta (Brunetti, 1910)
 australis Malloch (1932)
 spinigera Malloch (1932)
 schillingsi (Grunberg, 1906)
 minuta (Bezzi, 1892)
 longipalpis (Roubaud, 1906)
 potrix (Enderlein, 1928)
 meridiana Zumpt (1973)
 thirouxi thirouxi (Roubaud, 1906)
 hirudo (Enderlein, 1924)
 minuta (Enderlein, 1928, nec Bezzi)
 thirouxi potans (Bezzi, 1907)
 pallidipes (Roubaud, 1907)
 thirouxi titillans (Bezzi, 1907)
 irritans (Rondani, 1862, nec Linnaeus)
 tibialis (Hough, 1900, nec Robineau-Desvoidy)
 equina (Enderlein, 1928)
 latirostris (Enderlein, 1928)
 bovina (Peus, 1937)
 scolopax (Peus, 1937)

Supplement I:
Annotated Bibliography of the Horn Fly,
***Haematobia irritans irritans* (L.)**
Including References on the Buffalo Fly,
***H. irritans exigua* (de Meijere), and**
Other Species Belonging to the Genus *Haematobia*

Clyde E. Morgan and Gustave D. Thomas¹

This supplement is the first to "Annotated Bibliography of the Horn Fly, *Haematobia irritans* (L.), Including References on the Buffalo Fly, *H. exigua* (de Meijere), and Other Species Belonging to the Genus *Haematobia*," published in 1974 (U.S. Dep. Agric., Misc. Publ. No. 1278).

Included in this supplement are literature citations on *Haematobia* published from 1971 through 1975 and publications prior to 1971 omitted in the previous bibliography. Literature containing original information relating to biology, systematics, distribution, and control of *Haematobia* is annotated. Literature summarizing known information of minor importance is listed.

Abbreviations used in the citations are constructed according to the rules of the "American National Standard for Abbreviation of Titles of Periodicals" and can be found in "BIOSIS 1974 List of Serials with Coden, Title Abbreviations, New, Changed, and Ceased Titles," published by Bio-Science Information Service of Biological Abstracts.

In 1973, F. Zumpt, South Africa, presented a treatise revising the Stomoxyinae of the world. We list his classification of *Haematobia* to assemble the many synonymous species in this and the previous bibliography, and to bring information on the genus up to date. Currently *Haematobia* is comprised of six species including five subspecies.

¹Biological Control of Insects Research, Agricultural Research Service, Columbia, Mo.

REFERENCES

1. ABDULLAEV, U., and E. I. GAN. 1972. Pasture horn flies (*Haematobia* spp.). pp. 97-110. In Sultanov, M. A. (ed.) The Parasitic Arthropods of the Ferghana Valley. (In Russian.) Tashkent. Uzbek SSR, Izdatel'stvo Fan Uzbekskoi SSR. 152 pp.
2. ABLES, J. R., and M. SHEPARD. 1974. Hymenopterous parasitoids associated with poultry manure. Environ. Entomol. 3(5): 884-886.
3. ADERS, W. M. 1917. Insects injurious to man and stock in Zanzibar. Bull. Entomol. Res. 7: 399.
Lyperosia (= *Haematobia*) *minuta* showed preference for windy steppes.
4. ALDRICH, J. M. 1905. A catalog of North American Diptera. Smithson. Misc. Collect. 46, No. 1444, pp. 530-531.
5. ———. 1930. Notes on synonymy of Diptera, No. 4. Entomol. Soc. Wash. Proc. 32:28.
Horn fly, *Lyperosia* (= *Haematobia*) was believed synonymous with the moose fly, *Haematobia* (= *Haematobosca*).
6. ALICATA, J. E. 1969. Parasites of man and animals in Hawaii. Basel. pp. 62, 135.
H. i. irritans was listed as the intermediate host for *Stephanofilaria stilesi*.
7. ALLEN, W. C. 1973. Insects and other arthropods of importance during 1972 in houses, warehouses, and on livestock in Ontario, Canada. Entomol. Soc. Ont. Proc. 103: 2-3.
8. ANDERSON, J. R. 1974. Symposium on reproduction of arthropods of medical and veterinary importance. II. Meeting of the sexes. J. Med. Entomol. 11(1): 13-14.
9. ——— and J. B. Hov. 1972. Relationship between host attack rates and CO₂ baited insect flight trap catches of certain *Symphomyia* species. J. Med. Entomol. 9(5): 388.
10. ANONYMOUS. 1930. Principales plagas y enfermedades de los cultivos en la Republica Mexicana incluyendo las mas importantes de los Estados Unidos de Norte America. Talleres Graficos de la Secretaria de Agricultura y Fomento. Tacubaya, p. 243.
11. ———. 1971. Applied ecology used against a fly problem in Australia. Entomol. Rec. J. Var. 83(11): 353-354.
12. ———. 1972. Dung beetles on the move from rural research CSIRO. Wildl. Aust. 9(4): 101-104.

13. ——— 1973. Dust bags do control horn flies. N. Mex. Ext. News 53(3): 2-3.
14. ——— 1974a. Dung beetles combat flies. Agric. Res. (Washington, D.C.) 23(4): 7.
15. ——— 1974b. Australian meat research committee report. 8th Annu. Rep., year ended June 30, 1974. Australian Meat Board, Sydney. 144 pp.
16. ——— 1976. Trojan horse. Agric. Res. (Washington, D.C.) 24(7): 11.
17. ARDO, P. 1957. Studies in the marine shore dune ecosystem, with special reference to the dipterous fauna. Opusc. Entomol. Suppl. 14: 149.
18. ARTHUR, B. A. 1975. Self-oiling back-rubber for buffalo fly control. Queensl. Agric. J. 101(3): 373-377.
19. ASKEW, R. R. 1971. Parasitic insects. London, 316 pp.
20. ASS, M. YA. 1957. Diagnostic morphology of larval synanthropic flies. (In Russian.) Cesk. Parasitol. 4: 33-34, 48-49.
Discussed the spiracular plates of *H. i. irritans*, *Haematobosca stimulans*, and other larvae.
21. AUDCENT, H. 1932. Bristol insect fauna. Bristol Nat. Soc. Proc. (4) 7, pt. 5: 358-370.
22. AUSTEN, E. E. 1926. Concerning the nomenclature and synonymy of certain Oriental and East African bloodsucking muscids (Diptera). Bull. Entomol. Res. 16: 307.
23. AUSTRALIA COMMONWEALTH SCIENTIFIC AND INDUSTRIAL RESEARCH ORGANIZATION. 1970. Buffalo fly. Div. Entomol. Annu. Rep. 1969-70, p. 46.
24. ——— 1971. Dung beetles, buffalo fly. Div. Entomol. Annu. Rep. 1970-71, pp. 80-83, 96-97.
25. ——— 1972. Dung breeding flies. Div. Entomol. Annu. Rep. 1971-72, pp. 1, 72-73.
26. AYYAR, T. V. R. 1920. Note on a musciphagous wasp (*Bembex lunata*). Rep. 3d Entomol. Meet. Proc., Pusa, 1919. 3: 909-910.
B. lunata observed collecting *Stomoxys*, *Lyperosia* (= *Haematobia*), and *Philoematomyia* near a cow.
27. BABCOCK, O. G. 1920. Some common parasites of live stock in Colorado. Colo. Ext. Bull. Series 1, No. 166-A, pp. 6-8.

28. BADESCU, C. 1971. Contributions to the study of the dissemination and control of zoophilous flies on cattle breeding farms. (In Romanian.) Lucr. Stiint. Inst. Agron. 'N Balcescu' Bucur. Ser. C Med. Vet. 14: 271-277.

Haematobia was abundant at one of the three study farms.

29. BATTE, E. G. 1972. Differential diagnosis of parasitic dermatitis of cattle. J. Am. Vet. Med. Assoc. 161(11): 1265-1268.

30. BEADLES, M. L., J. A. MILLER, W. F. CHAMBERLAIN, and others. 1975. The horn fly: methoprene in drinking water of cattle for control. J. Econ. Entomol. 68(6): 781-785.

Devices were constructed to meter 0.34 mg of methoprene/liter of water automatically into the troughs of drinking water of a 350-animal herd of cattle. Development of horn flies in manure was completely inhibited. The number of flies was reduced from an average of 360 to 7 per animal.

31. BENBROOK, E. A. 1946. List of parasites of domestic animals of North America. Minneapolis, p. 41.

32. BENOIT, P. L. G. 1964. Mission de zoologie medical au Maniema (Congo, Leopoldville) (P.L.G. Benoit, 1959). 14. Diptera-Stomoxidae. Ann. Mus. R. Afr. Cent., Tervuren (Ser. 8°). Sci. Zool. No. 132, pp. 265-268.

H. minuta and *H. t. thirouxi* were present in the Congo.

33. BERRY, I. L. 1973. Improved system for measuring flying activity of insects by detecting static charges. J. Econ. Entomol. 66(3): 820-822.

H. i. irritans was one of the insects used for testing static charge system.

34. ——— R. L. HARRIS, J. L. ESCHLE, and J. A. MILLER. 1974. Large scale rearing of horn flies on cattle. U.S. Dep. Agric., Agric. Res. Serv., ARS (Ser) South. Reg. S-35. 6 pp.

Adult horn flies were maintained on steers and allowed to oviposit directly on the feces from the steers. The egg-bearing feces were combined with an artificial larval-rearing medium which increased the survival and size of the resulting pupae. Procedures developed during the study resulted in an average daily production of 24,800 pupae from about 6,500 flies on each steer.

35. ——— J. A. MILLER, C. D. SCHMIDT, and J. L. ESCHLE. 1973. Simulation of horn fly populations. Trans. ASAE (Am. Soc. Agric. Eng.) 16(3): 482-485.

A deterministic model of population growth was developed and used for describing growth of horn flies in the laboratory.

36. BEZZI, M. 1917. Studies in Philippine Diptera, II. Philipp. J. Sci., Sect. D. 12(3): 107.
H. i. exigua was found on Luzon.
37. BISHOPP, F. C. 1913. The stable fly (*Stomoxys calcitrans* L.), an important live stock pest. J. Econ. Entomol. 6: 112-126.
38. ——— 1952. Insect friends of man. U.S. Dep. Agric. Yearb. 1952: 82.
39. BLAGOVESHCHENSKY, D. I. 1937. Materials pertaining to the fauna of the external parasites (Arthropoda) of animals of Kazalinsk and some other districts of southern Kazakhstan. (In Russian.) Tr. Akad. Nauk Kazakhskoi SSR (Alma-ata). 2: 84 pp.
H. i. irritans was present in Kazakhstan.
40. Blume, R. R. 1972. Additional insects associated with bovine droppings in Kerr and Bexar counties, Texas. J. Econ. Entomol. 65(2): 621.
41. ——— J. J. MATTER, and J. L. ESCHLE. 1973a. Biting flies (Diptera: Muscidae) on horses: laboratory evaluation of 5 insecticides for control. J. Med. Entomol. 10(6): 596-598.
H. i. irritans was controlled with applications of crotoxyphos, coumaphos, carbaryl, malathion, and pyrethrins.
42. ——— J. J. MATTER, and J. L. ESCHLE. 1973b. *Onthophagus gazella*: effect on survival of horn flies in the laboratory. Environ. Entomol. 2(5): 811-813.
Varying pairs of *O. gazella* were exposed to varying quantities of cow manure containing eggs or larvae of *H. i. irritans*. Control of emerging adult flies ranged from 0 to 98.4 percent.
43. BOHART, G. E., and J. L. GRESSITT. 1951. Filth-inhabiting flies of Guam. Bull. Bernice P. Bishop Mus. No. 204: 13, 21-22, 29, 35, 47, 50, 54, 58, 117-119.
Taxonomy of *Haematobia* and related genera on Guam was presented.
44. BORNEMISSZA, G. F. 1970. Insectary studies on the control of dung breeding flies by the activity of the dung beetle, *Onthophagus gazella* F. (Cole: Scarabaeinae). J. Aust. Entomol. Soc. 9: 31-41.
45. BORNEMISSZA, I. 1971. Biologische regelung der bevölkerungsdichte von *Haematobia exigua* durch ansiedlung von dungkafern in Australien. Anz. Schaedlingskd. 44: 9-10.
46. BOURNE, J. R., and K. L. HAYS. 1973a. Ration fed to cows affects horn flies. Ala. Agric. Exp. Stn., Highlights Agric. Res. 20(1).

47. ——— and K. L. HAYS. 1973b. Predators of the horn fly in the Piedmont of Alabama. Ala. Agric. Exp. Stn., Highlights Agric. Res. 20(3) : 15.
48. BRAUER, F., and J. E. BERGENSTAMM. 1889. Die zweiflugler des Kaiserlichen Museums zu Wien. IV. Vorarbeiten zu einer monographie der muscaria schizometopa. Pars I. Akad. Wiss. Math-Naturwiss. Kl. 56: 155.
49. BRAUNS, A. 1959. Autokologische untersuchungen uber die thal-assicolen zweiflugler (Diptera) im schlesewig-holsteinischen bereich der Nord-und Ostsee. Arch. Hydrobiol. 55: 545.
50. BRIMLEY, C. S. 1938. The insects of North Carolina. Raleigh, p. 372.
51. BRITTON, W. E. 1920. Check list of the insects of Connecticut. Conn. State Geol. Nat. Hist. Surv. Bull. 31: 197.
52. BROWN, R. S. 1950. DDT content of milk as a result of cow and barn spraying. 74th Annu. Rep. Ont. Agric. Coll. Exp. Farm, 1949, p. 50.
53. BRUNETTI, E. 1917. Diptera of the Simla District. Rec. Indian Mus. 13: 59-101.
54. BUEI, K. 1952. On the stable fly (*Stomoxys*) and the horn fly (*Lyperosia*) of Osaka prefecture. Jpn. J. Sanit. Zool. 3(1-2): 20-25.
Discussed morphology of the flies including differences in antennae, palps, hind harsi, phalosomes, and parameres.
55. BURGER, J. F., and J. R. ANDERSON. 1974. Taxonomy and life history of the moose fly, *Haematobosca alcis*, and its association with the moose, *Alces alces shirasi* in Yellowstone National Park. Ann. Entomol. Soc. Am. 67(2) : 204-214.
56. BURNETT, E. M. 1970. Pesticide residues in fat of cattle treated with backrubbers impregnated with coumaphos, methoxychlor, or DDT. Pestic. Sci. 1(2) : 70-72.
57. BURNS, E. C., G. R. MCCOY, D. G. MELANCON, and others. 1975. Effects of horn flies on rate of gain of stocker beef cattle. Lives. Prod. Day, La. State Univ., Anim. Sci. Dep. 15: 258-261.
Rate of gain was reduced when fly populations exceeded 200/cow.
58. ——— D. G. MELANCON, J. C. CARPENTER, and J. I. FEAZEL. 1975. Studies with Altosid IGR in mineral formulations for the control of horn flies on beef cattle. Lives. Prod. Day, La. State Univ., Anim. Sci. Dep. 15: 262-270.
Fly populations were reduced on cattle and emergence of flies from manure was eliminated.

59. BURSSELL, E. 1975. Substrates of oxidative metabolism in dipteran flight muscle. *Comp. Biochem. Physiol. B. Comp. Biochem.* 52(2): 235-238.

A comparison was made of the capacity of mitochondria from the flight muscles of various Diptera including *Haematobia* sp. to oxidize pyruvate, x-glycerophosphate, and proline.

60. BUTLER, J. F., and N. I. GREER. 1973. Toxicity of SD-8447 and dichlorvos to larvae of the horn fly, *Haematobia irritans* (Diptera: Muscidae) in manure of insecticide-fed cattle. *Fla. Entomol.* 56(2): 103-105.

When used as feed additives, both insecticides caused significant mortality of larvae in manure samples from cattle.

61. BUXTON, P. A. 1955. The natural history of tsetse flies. London School of Hygiene and Tropical Medicine, Mem. No. 10, pp. 738-739.

62. CAMPION, D. G. 1972. Insect chemosterilants: a review. *Bull. Entomol. Res.* 61: 633.

63. CANNON, F. M. 1951. Insects affecting man and animals. *Can. Insect Pest Rev.* 29: 105.

64. CARPENTER, G. H. 1899. Insects, their structure and life. London, p. 265.

65. CATES, S. 1938. Branching road of science. *Country Gentleman* 108(9): 23, 67.

66. CHAMBERLAIN, W. F. 1974. Horn flies: emergence, survival, and sterility after cobalt-60 irradiation of pupae. *J. Econ. Entomol.* 67(3): 381-383.

Irradiation of colony-reared 30-hr-old pupae with 2,500 rad completely inhibited fly emergence; with irradiation of 54-hr-old pupae, some flies emerged, but all died within 3 days. Adults from irradiated 78-hr-old pupae emerged, but unusual numbers died within 3 days. Neither emergence nor survival was affected adversely when 102 or 126-hr-pupae were irradiated. Pupal irradiation with 2,000 and 2,500 rad induced sterility in the adult flies.

67. CHOW, C. Y., T. C. HUANG, and T. F. YUE. 1950. Bibliography of Chinese arthropods of medical and veterinary importance. *Q.J. Taiwan Mus. (Taipei)* 3 (3): 157-185.

68. Christensen, C. M. 1975. The effect of testosterone propionate on the attractiveness of cattle to horn flies. *North Cent. Branch Entomol. Soc. Am. Proc.* 30: 97.

Testosterone propionate treated steers and bulls had larger sebaceous gland cells than the control steers. *Inspection of 1973*

daily horn fly counts indicated that bulls were the most attractive. Treated steers were more attractive than control steers 6 to 9 days after treatment. Regression analysis indicated a positive relationship between horn fly attractance and sebaceous gland parameters.

69. COCKRILL, W. R. 1974. The buffaloes of Latin America. pp. 676-707. In Cockrill, W. R. (ed.) 1974. The Husbandry and Health of the Domestic Buffalo. FAO, Rome. 993 pp.
Buffalo flies (*Haematobia*) did not appear to exist in Brazil.
70. COLLINS, J. E., and C. J. WAINWRIGHT. 1934. Some Diptera collected in the south of England in 1930-33. J. Soc. Brit. Entomol. 1(1): 21.
H. i. irritans was found in Arne, Dorset.
71. COWSERT, W. C., J. W. SCALES, and J. W. WARD. 1945. DDT starts work for dairy farmers. Miss. Farm Res. 8(12): 1-2.
72. CREELMAN, I. S. (ed.) 1968. Horn fly. Can. Agric. Insect Pest Rev. 46: 25.
73. ——— 1969. Horn fly. Can. Agric. Insect Pest Rev. 47: 32.
74. ——— 1970. Horn fly. Can. Agric. Insect Pest Rev. 48: 34.
75. ——— 1971. Horn fly. Can. Agric. Insect Pest Rev. 49: 38.
76. ——— 1972. Horn fly. Can. Agric. Insect Pest Rev. 50: 35.
77. CRESS, D. C., and T. R. GREATHOUSE. 1972. Evaluation of different methods of controlling face and horn flies on beef cow herds. Mich. Agric. Exp. Stn., Res. Rep. No. 166: 81-82.
Horn flies were controlled with 1 percent coumaphos dust bags; partial control was obtained with 7 percent phenothiazine salt blocks.
78. CRYSTAL, M. M. 1967a. Carbon dioxide anesthesia of untreated and chemosterilant-treated screwworm flies, *Cochliomyia hominivorax* (Coquerel). J. Med. Entomol. 4(4): 418.
79. ——— 1967b. Reproductive behavior of the laboratory-reared screwworm flies. (Diptera: Calliphoridae). J. Med. Entomol. 4(4): 445.
80. CZERNY, L., and G. STROBL. 1909. Spanische dipteren. III. Beitrag. Vehr. Zool.-Bot. Ges. Wien 59: 235.
H. i. irritans was collected in Spain.
81. DALRYMPLE, W. H. 1913. Some of the more important insects affecting our farm animals. Am. Vet. Rev. 42: 424-427.
82. D'ASSIS FONSECA, E. C. M. 1968. Diptera: Cyclorrhapha; Calyptrata. Section (b) Muscidae. Handbooks for the Identification of British Insects 10: Pt. 4 (b), p. 12.

83. DEL FOSSE, E. S., and E. U. BALSBAUGH. 1973. Side effects of ULV organophosphates, aerially applied for fly control on cattle, on dung beetle fauna. North Cent. Branch Entomol. Soc. Am. Proc. 28: 195.
No significant effects to the dung biota could be attributed to ULV applied Ravap or malathion. ULV application of stirofos diminished several groups of dung beetles.
84. ——— and E. U. BALSBAUGH. 1974. Effects of ULV organophosphates on horn flies and face flies of cattle, and on bovine coprocenosis. Environ. Entomol. 3(6): 919-922.
Same as above.
85. DETHIER, V. G. 1956. Repellents. Annu. Rev. Entomol. 1: 187.
86. DERBENEVA-UKHOVA, V. P. 1942. On the development of ovaries and on the imaginal nutrition in dung flies. (In Russian.) Med. Parazitol. 11(4): 85-97. Abstr. in Rev. Appl. Entomol. 1944, 32: 118.
87. ——— 1952. Flies and their epidemiological importance. (In Russian.) Moscow, 272 pp.
88. DOBREANU, E., A. BERTEANU, and A. DUMITREASA. 1962. Determinator al Mustelor Sinantrope din R.P.R. Editura Academiei Republicii Populare Romine. Bucuresti, pp. 68-71.
89. DOVE, W. E. 1945. Summary of DDT experiments on insects that affect man and animals. U.S. Dep. Agric., Entomol. Plant Quarantine. E-673, pp. 5, 6.
90. DOWNES, J. A. 1958. The feeding habits of biting flies and their significance in classification. Annu. Rev. Entomol. 3: 260.
91. DRABER-MONKO, A. 1966. Eine neue art der gattung *Siphona* Meig. (Diptera: Muscidae) aus der Mongolei. Bull. Acad. Pol. Sci., Ser. Sci. Biol. 14(4): 225-232.
Described *Siphona ryszardi* (= *Haematobosca stimulans*).
92. DRAKE-BROCKMAN, R. E. 1910. Notes on the bloodsucking Diptera met with in eastern and south-eastern Abyssinia. Bull. Entomol. Res. 1: 55.
Lyperosia (= *Haematobia*) *minuta* was common near Mt. Fantali.
93. DRENSKI, P. 1957. Die familie der stechfliegen. (In Bulgarian.) (German summary.) Izvestiya na Zoologicheskaya Institut B'lgarska Akademiya na Naukite 6: 449-450.
Description of stomoxiine flies found in Bulgaria.

94. DROBISHCHENKO, N. I., and V. A. SHOL'. 1973. The infection of stable fly, *Haematobia stimulans*, with larvae of *Setaria cervi*. Deer nematodes, insect vectors. (In Russian.) Izv. Acad. Nauk. Kaz. SSR Ser. Biol. Nauk. 5: 20-24.
95. ———. 1975. The developmental cycle of *Haematobia stimulans* Mg. (Diptera: Muscidae), an intermediate host of the nematode *Setaria cervi* Rud., in the Southern Altai. (In Russian.) Entomol. Obozr. 54(2): 342-345.
96. DULAC, A. 1960. Etude des Anthomyidae de la collection E. Hesse (premiere note). Travaux du Laboratoire Zoologie Stat. Aquicole Grimaldi de la Faculte des Sci. Dijon 35: 19.
97. DUREN, E., and L. E. O'KEEFE. 1972. Horn fly control with dust bags: effect on weight gains. West. Section, Am. Soc. Anim. Sci. Proc. 23: 53-56.
Weight gains were recorded for treated cattle.
98. EASTAWAY, B. 1974. The buffalo fly in Queensland. Queensl. Agric. J. 100(6): 221-224.
99. EDDY, G. W. 1952. Flies on livestock. U.S. Dep. Agric. Yearb. 1952: 657-661.
100. EDWARDS, F., H. OLDROYD, and J. SMART. 1939. British blood-sucking flies. (British Museum) London, pp. 115-118.
101. EMDEN, F. I. VAN. 1948. 14. Diptera: Muscidae. British Museum Expedition to S.W. Arabia 1: 174.
Lyperosia (= *Haematobia*) *minuta* was recovered from camels in Western Aden Protectorate.
102. ———. 1965. Diptera, vol. 7, Muscidae, part 1. In The Fauna of India and Adjacent Countries. Zoological Survey of India, pp. 175-178.
103. ENDERLEIN, G. 1924. Eine neue blutsaugende Muscidae aus Afrika. Konowia 3: 51-52.
Described *Haphospatha hirudo* (= *H. t. thirouxi*).
104. ESCHLE, J. L., S. E. KUNZ, C. D. SCHMIDT, and others. 1973. Suppression of a population of horn flies with the sterile-male technique. Environ. Entomol. 2(6): 976-980.
Sterilized *H. i. irritans* were released into a semi-isolated population in West Texas. Initial releases caused a downward trend in reproduction. Continued releases caused a 98 percent decrease in reproduction during the last 3 weeks of the 16-week study, and control exceeded 70 percent for 10 weeks. When the sterile/fertile ratio dropped below 6:1, the decrease in reproduction dropped below 90 percent.
105. ——— and A. MILLER. 1968. ULV application of insecticides to cattle for control of the horn fly. J. Econ. Entomol. 61(6): 1617-21.

- A sprayer was developed to provide animal-activated ULV application of insecticides to cattle. Fly control was obtained with all of the insecticides tested.
106. ——— D. D. OEHLER, B. F. HOGAN, and M. C. IVEY. 1973. Coumaphos on cable type backrubbers: control of horn flies and determination of residues in milk of dairy cattle. *J. Econ. Entomol.* 66(1): 290-291.
One to two percent solution of coumaphos gave control of horn flies throughout the 4-week test. No residues were detected in milk.
 107. ——— C. D. SCHMIDT, and J. J. MATTER. 1973. Horn fly: preliminary evaluation of media for rearing larvae. *Ann. Entomol. Soc. Am.* 66(2): 352-354.
H. i. irritans were reared on bovine feces and artificial media containing various proportions of sugar cane bagasse, whole wheat flour, fishmeal, baking soda, and water. Bovine feces appeared essential for maximum fly production, but an artificial medium could be substituted for some of the feces.
 108. ESSIG, E. O. 1926. *Insects of Western North America*. New York, p. 590.
 109. ESSIG, H. W., W. A. PUND, and C. F. BAILEY. 1964. A study of horn fly control in beef cattle. *Miss. State Univ. Agric. Exp. Stn. Inform. Sheet* 859, 2 pp.
Mineral blocks containing ronnel were used for horn fly control.
 110. FABRICIUS, J. C. 1805. *Systema Antliatorum secundum ordines, genera, species*. *Brunsvigae* (=Brunswick), p. 281.
 111. FAN, C. T. 1965. Key to the common synanthropic flies in China. (In Chinese.) *Academy of Sciences, China*, 330 pp.
We have not seen this reference, but it is listed in Shinonaga and Kano, 1971, and Zumpt, 1973.
 112. FAY, R. W., and J. W. KILPATRICK. 1958. Insecticides for control of adult Diptera. *Annu. Rev. Entomol.* 3: 401-420.
 113. FERNANDEZ-SOUSA, J. M., J. G. GAVILANES, A. M. MUNICO, and others. 1975. Primary structure of cytochrome c from the insect *Ceratitis capitata*. *Biochem. Biophys. Acta* 393(2): 358-367.
Amino acid sequence of cytochrome c from *C. capitata* was compared with that of *H. i. irritans*.
 114. FERRAR, P. 1973. The CSIRO dung beetle project. *Wool Technol. Sheep Breed.* 20(1): 73-75.
 115. ——— 1975. Disintegration of dung pads in north Queensland before the introduction of exotic dung beetles. *Aust. J. Exp. Agric. Anim. Husb.* 15(74): 325-329.

116. FINCHER, G. T., R. DAVIS, and T. B. STEWART. 1971. Flight activity of coprophagous beetles on a swine pasture. *Ann. Entomol. Soc. Am.* 64(4): 855-860.
117. FORDHAM, W. J. 1945. A preliminary list of the Diptera of Northumberland and Durham (excluding the Cecidomyiidae). *Trans. Nat. Hist. Soc. Northumberl. Durham Newcastle-Upon-Tyne (New Ser.)* 7: 197-265.

Haematobia (= *Haematobosca*) *stimulans* was included in the Diptera list.
118. FREY, R. 1924. Diptera Brachycera. In Dampf, A. *Zur Kenntnis der estlandischen Hochmoorfauna. Sitzungsberichte Naturforscher-Gesellschaft Universitat Dorpat* 31: 55.

Haematobia (= *Haematobosca*) *stimulans* was listed.
119. GAMAL-EDDIN, F. M. 1968. Field studies on the biting cycle of two bloodsucking flies (*Siphona irritans* and *Musca crassirostris*) in Egypt. *J. Egypt Vet. Med. Assoc.* 28(1/2): 83-91.

On cattle, *H. i. irritans* has two peaks of biting activity during the different seasons—during spring and summer the afternoon peaks were slightly higher than the morning peaks; the reverse was true during late autumn and winter. On donkeys, only one peak of biting activity was noted (in the morning).
120. ——— 1971a. Field and laboratory studies on the life cycle of the blood-sucking, ectoparasitic fly *Siphona irritans*, Lin. in Egypt (Diptera: Muscidae). *J. Egypt. Vet. Med. Assoc.* 31(3/4): 243-250.

Oviposition lasted an average of 9.5 days at 30° C and 13 days at 25° C. At natural temperatures, the average number of eggs deposited was at its lowest (20) for the year in October and rose gradually to a maximum of 125 in May, after which it again declined. At constant temperatures of 20° and 30° C, an average of 74.3 and 142.5 eggs were deposited respectively.
121. ——— 1971b. Studies on the behavior of the blood-sucking fly *Siphona irritans* Lin. towards some environmental factors, to pave the way for proper control. A. The adult stage. (2). Reactions to temperature. *J. Egypt. Vet. Med. Assoc.* 31(3/4): 251-262.

Lab studies showed that in choice-chamber experiments, *H. i. irritans* exhibited a preference for temperatures between 20° and 35° C. Preconditioning had no effect on the subsequent behavior of the flies.
122. ——— 1972a. Field and laboratory studies on the life cycle of the blood-sucking ectoparasitic fly *Siphona irritans*, Lin. in Egypt. (2) The egg stage. *J. Egypt. Vet. Med. Assoc.* 32 (1/2): 45-53.

Duration of the egg stage varied from an average of 30 hr at 20° C to 11.2 days at 33.5° C. Eggs held below 90 percent RH did not hatch. As the temperature increased, the lower limit of humidity at which hatching occurred was reduced.

123. ——— 1972b. Field and laboratory studies on the life cycle of the blood-sucking ectoparasitic fly *Siphona irritans*, Lin. in Egypt. 3. The larval and pupal stages and their micro habitats in nature. J. Egypt. Vet. Med. Assoc. 32(1/2): 55-74.

Duration of the three larval instars varied from 13 days at 18° C to 3.8 days at 32° C; duration of the pupal stage varied from 13.5 days at 18° C to 3.9 days at 31° C. Pupal period was shorter at 98 percent RH than at 90, 95, or 100 percent RH at 25° C. Field studies included observations on temperature, RH, and pH of cow pats. In Egypt optimum developmental temperatures occurred in the spring.

124. ——— 1972c. Studies on the behavior of the blood-sucking fly *Siphona irritans*, Lin. toward some environmental factors, to pave the way for proper control. (Diptera: Muscidae) A. The adult stage. (3) Smell reaction and olfactory receptors. J. Egypt. Vet. Med. Assoc. 32 (3/4): 229-246.

Reactions of *H. i. irritans* to the smell of animals and dung were studied. Various combinations of smell, temperature, and humidity were tested. The degree of moistening of the host's body was found to determine the degree of abundance of flies on the host.

125. GJULLIN, C. M., and V. D. BEVILL. 1972. Insect chilling table. J. Med. Entomol. 9(3): 266.

126. GLASER, R. W. 1923. The effect of food on longevity and reproduction in flies. J. Exp. Zool. 38(3): 383-412.

H. i. irritans could be kept in captivity for 2 to 25 days but ceased egg laying 2 to 5 days after capture.

127. GMELIN, J. F. 1790. Caroli a Linne', systema naturae per Regna Tria Naturae. (ed. 13). vol. 1: Regnum Animale, pt. 5. Lipsiae (=Leipzig). p. 2892.

128. GOBLE, H. W. 1948. Insects attacking man and domestic animals. Can. Insect Pest Rev. 26: 56, 80, 85.

129. ——— 1950. Insects affecting man and domestic animals. Can. Insect Pest Rev. 28: 56, 92.

130. GONZALEZ, R. H. 1968. *Haematobia irritans* (L.) en Chile. Rev. Chil. Entomol. 6: 142.

Collected *H. i. irritans*, not previously known in Chile.

131. GONZALEZ-RINCONES, R., and L. GUYON. 1953. Classificacion general de los Dipteros. Caracas, pp. 118-125.

132. GOTWALD, W. H. 1965. A checklist and keys to the Muscinae and Stomoxydinae (Diptera: Muscidae) of Pennsylvania. *Entomol. News* 76: 199-210.
133. GRAHAM, O. H., R. O. DRUMMOND, and R. A. HOFFMAN. 1968. Possibilities of the sterile-male technique for the control of livestock insects in the United States of America. *In* Control of Livestock Insect Pests by the Sterile-Male Technique. (Panel Proc. Ser.) Int. At. Energy Agency. pp. 41-44.
134. GREENBURG, B., and D. POVOLNY'. 1971. Bionomics of flies. *In* Greenburg, B. Flies and Disease. vol. 1. Princeton p. 65.
135. GREGOR, F., and D. POVOLNY'. 1958. Versuch einer klassifikation der synanthropen fliegen. *J. Hyg. Epidemiol. Microbiol. Immunol.* (Prague) 2: 205-216.
136. ——— and D. POVOLNY'. 1960. Beitrag zur kenntis der synanthropen fliegen ungarns. *Cas. Cesk. Spolecnosti Entomol.* 57(2): 158-177.
137. GRIFFITHS, G. C. D. 1972. The Phylogenetic classification of Diptera Cyclorrhapha. The Hague, pp. 146-147.
Abdominal structures were used in the classification of *H. i. irritans*.
138. GRIFFITHS, R. B. 1974. Parasites and parasitic diseases. pp. 236-275. *In* Cockrill, W. R. (ed.) The Husbandry and Health of the Domestic Buffalo. FAO, Rome, 993 pp.
H. i. exigua populations on buffalo were larger than those on cattle; buffalo appeared not to suffer from feeding activity of the flies.
139. GRUNBERG, K. 1907. Die Blutsaugenden Dipteren. Jena, pp. 160-161.
140. HAESELBARTH, E., J. SEGERMAN, and F. ZUMPT (ed.). 1966. The arthropod parasites of vertebrates in Africa south of the Sahara (Ethiopian region). Vol. III, Publ. S. Afr. Inst. Med. Res. 13(52): 35-37.
141. HAEUSSLER, G. J. 1952. Losses caused by insects. U.S. Dep. Agric., Yearb. 1952: 144.
142. HAFEZ, M., and F. M. GAMAL-EDDIN. 1968. The feeding habits of the horn fly (*Siphona irritans*) with special reference to its biting cycle in nature in Egypt (Diptera: Muscidae). *Bull. Soc. Entomol. Egypte* 50(59): 41-52.
143. HARDY, D. E. 1952. Additions and corrections to Bryan's check list of the Hawaiian Diptera. *Hawaii. Entomol. Soc. Proc.* 14: 480-481.

144. HARLEY, J. M. B. 1965. Seasonal abundance and diurnal variations in activity of some *Stomoxys* and Tabanidae in Uganda. Bull. Entomol. Res. 56(2): 319-332.
145. HARRIS, R. L., W. F. CHAMBERLAIN, and E. D. FRAZAR. 1974. Horn flies and stable flies: free choice feeding of methoprene mineral blocks to cattle for control. J. Econ. Entomol. 67(3): 384-386.
- With 0.94, 0.12, and 0.01 percent methoprene in mineral blocks, development of *H. i. irritans* in the field was inhibited 87 percent. Bioassay data in the laboratory indicated better than 97 percent inhibition.
146. ——— E. D. FRAZAR, and R. L. YOUNGER. 1973. Horn flies, stable flies, and house flies: development in feces of bovines treated orally with juvenile hormone analogues. J. Econ. Entomol. 66(5): 1099-1102.
- H. i. irritans* development was inhibited in feces of cattle treated orally with 1 g/day of Hoffman-La Roche Ro7-9767 and 0.7 mg/day of Zoecon ZR-515.
147. ——— J. A. MILLER, and E. D. FRAZAR. 1974. Horn flies and stable flies: feeding activity. Ann. Entomol. Soc. Am. 67(6): 891-894.
- An electronic bitometer was used to determine the feeding activity of *H. i. irritans* feeding on a cow. Females spent an average of 163 min/day in feeding and averaged 38.4 feedings/day. Males averaged 24 feedings/day and spent an average of 96 min/day in feeding. In the laboratory female horn flies averaged 10 feedings/day and spent an average of 15.3 min/day in feeding.
148. HASEGAWA, T., H. HAYAKAWA, and T. CHIBA. 1973. Sand flies in Tohoku area, Japan. (In Japanese.) Jpn. J. Sanit. Zool. 23(4): 271.
149. HAUFÉ, W. O. 1970. An economic evaluation of horn fly control on yearling cattle treated with DDT and coumaphos. Annu. Meeting Can. Soc. Anim. Production Proc., July 5-9, Ottawa. pp. 48-49.
150. ——— 1973a. Horn fly. Can. Agric. Insect Pest Rev. 51: 48.
151. ——— 1973b. Interaction of pesticidal toxicity, parasites, and reversible anticholinesterase activity as stresses on growth rate in cattle infested with horn flies (*Haematobia irritans*). Toxicol. Appl. Pharmacol. 25(1): 130-144.
- Differential rates of gain in groups of immature cattle exposed to *H. i. irritans* attack were used to distinguish the interaction of pesticidal activity and nonpesticidal pharmacologically induced stress with host-parasite reaction.

152. ——— 1974. Haematophagus flies—a component in the impact of environment on the health and productivity of cattle. Int. Livestock Symp. Proc., Am. Soc. Agric. Engin., Ser. Publ. SP-0174. pp. 203-208.
153. HAWKING, F., and M. WORMS. 1961. Transmission of filaroid nematodes. Annu. Rev. Entomol. 6:415.
154. HAYES, B. W., M. J. JANES, and D. W. BEARDSLEY. 1972. Dust bag treatments in improved pastures to control horn flies and cattle grubs. J. Econ. Entomol. 65(5): 1368-1371.
The 3-year study included various combinations of treatments of 1 and 5 percent coumaphos, 5 and 10 percent prolate, and 10 percent methoxychlor. All were effective in controlling *H. i. irritans* on Brahman-Angus or Brahman-Hereford cross-bred steers.
155. HEDLUND, L., L. HAHN, G. THOMAS, and H. D. JOHNSON. 1972. Effect of flies on body movement of dairy cows. (Abstract) J. Anim. Sci. 35: 166.
156. HEIMPEL, A. M. 1967. A critical review of *Bacillus thuringiensis* var. *thuringiensis* Berliner and other crystalliferous bacteria. Annu. Rev. Entomol. 12: 296.
157. HENNEGUY, L. F. 1904. Les insectes. Paris, p. 200.
158. HENNIG, W. 1952. Dipteren von den Suda-Inseln. IV. Fam. Muscidae. Beitr. Entomol. 2:93.
159. ——— 1964. Fam. 63b. Muscidae lfg. 249: 1009-1056. In Lindner, E. (ed.), Die Fliegen der Palaearktischen Region. Stuttgart.
160. ——— 1965. Vorarbeiten zu einem phylogenetischen system der Muscidae (Diptera: Cyclorrhapha). Stuttg. Beitr. Naturkd. 141: 76-79.
161. ——— 1968. Die Larvenformen der Dipteren. 3. Teil. Berlin pp. 365-366, 372-373.
162. HERTING, B. 1957. Das weibliche postabdomen der calyptraten fliegen (Diptera) und sein merkmalswert fur die systematik der gruppe. Z. Morph. Okol. Tiere 45: 436.
163. HINTON, H. E. 1960. The chorionic plastron and its role in the eggs of the Muscinae (Diptera). Q. J. Microsc. Sci 101: 313-332.
164. HOELSCHER, C. E., and R. L. COMBS. 1970. The horn fly. II. Comparative physiological studies of reproductive and diapaused pupae. Physiol. Zool. 43(4): 241-248.

Respiration rate decreased threefold to fivefold when diapaused pupae were compared with reproductive specimens. Moisture content was 63.7 percent for reproductive pupae and 44.7 percent for diapaused pupae. No differences observed in lipid content in pupal samples. Glycerol was not found in reproductive pupae, but was present in diapausing pupae.

165. ——— and R. L. COMBS. 1971. The horn fly. III. Sex ratio and factors affecting adult emergence. *Ann. Entomol. Soc. Am.* 64(4): 912-919.

A 1:1.35 male to female ratio was found from the total of 65,702 flies collected from pastured cattle during 2 years. A higher proportion of male horn flies was found in collections from the legs of calves. Temperature controlled emergence completely, but the effects of temperature were altered by photoperiod. Female flies emerged earlier than male flies in various tests. Photoperiod had the most pronounced effect on emergence when the larval and pupal stages were reared at a 12-hour reciprocal photophase of field conditions.

166. HOMAN, H. W., and E. P. DUREN. 1975. Horn fly control on range and pastures. Idaho Univ. Coop. Ext. Serv., Current Inform. Ser. No. 282, 2 pp.

167. HORI, K. 1960. Comparative anatomy of the internal organs of the calyptrate muscoid flies. I. Male internal sexual organs of the adult flies. *Sci. Rep. Kanazawa Univ.* 7(1): 34.

H. i. exigua was used in the study.

168. HORNE, W. T., E. O. ESSIG, and W. B. HERMS. 1923. Plant disease and pest control. Univ. Calif. Agric. Exp. Stn. Circ. 265, p. 63.

169. HORNING, B. 1959. Fliegen als ubertrager parasitarer wurmer. *Z. Angew. Zool.* 46: 338-342.

H. i. exigua was thought to be a vector of habronemiasis.

170. HOUGH, G. N. 1899. Some Muscinae of North America. *Biol. Bull. (Lancaster)* 1(1): 19-33.

Described differences between *H. serrata* (= *H. i. irritans*) and *H. alcis* (= *Haematobosca stimulans*).

171. ——— 1900. Notes on some European species of Calliphorinae, Muscinae Muscaeformes and Muscinae Ariciaeformes in the collection of the Hungarian National Museum. *Termesztudajzi Fuzetek.* 23: 249-250.

Discussed synonymy of *H. i. irritans*.

172. HU, CHING-FU. 1940. *Catalogus insectorum sinensium*. vol. 5. Peiping, pp. 341, 346-347.

173. HUKUSIMA, S., and S. TAKEDA. 1973. Biological studies of horse fly and horn fly in the Iwai pasture, Gifu prefecture located at upland region and repellent tests against them. (In Japanese.) Res. Bull. Fac. Agric. Gifu Univ. 34: 71-88.
174. HUCKETT, H. C. 1965. Family Muscidae. pp. 869-915. In Stone, A., C. W. Sabrosky, W. W. Wirth, R. H. Foote, and J. R. Coulson. (ed.) A Catalog of the Diptera of America North of Mexico. U.S. Dep. Agric., Agric. Res. Serv., Agric. Handb. No. 276, 1,696 pp.
175. ISHIJIMA, H. 1967. Revision of the third stage larvae of synanthropic flies of Japan (Diptera: Anthomyiidae, Muscidae, Calliphoridae, and Sarcophagidae). Jpn. J. Sanit. Zool. 18: 47-100.
Included descriptions of *Lyperosia exigua* (= *H. i. exigua*) and *H. sanguinolenta* (= *Haematobosca sanguinolenta*).
176. JACOBS, S. N. A. 1971. Applied ecology used against a fly pest problem in Australia. Entomol. Rec. J. Var. 83(11): 353-355.
General review of dung beetle program.
177. JANES, M. J., B. W. HAYES, and D. W. BEARDSLEY. 1970. Dust bags control horn flies in improved pastures. Sunshine State Agric. Res. Rep. 15(2/3): 15-16.
178. JONES, B. F. 1973. Control profit-stealing horn flies. Arkansas Cattle. Bus. 9(6): 7-9.
179. ——— and G. BARNES. 1974. Protect beef cattle from external parasites. Univ. Arkansas Coop. Ext. Serv. Leaflet 285 (Rev. Annu.) 6 pp.
180. KABOS, W. J. 1951. De Diptera Brachycera van het eiland Texel, oecologisch beschouwd. Tijdschr. Entomol. 93: 127.
181. ——— 1964. Tweevleugelige insekten—Diptera. VIII. De Nederlandse vliegen (Muscidae). Wet. Meded. K.N.N.V. (K. Ned. Natuurhist. Ver.) 53: 1-32.
H. (= *Haematobosca*) *stimulans* and *Lyperosia irritans* (= *H. i. irritans*) were among species discussed.
182. ——— 1975. Two-winged insects—Diptera. Flies of the Netherlands (Muscidae). 2d amplified edition. (In Dutch.) Wet. Meded. K.N.N.V. (K. Ned. Natuurhist. Ver.) No. 110, 64 pp.
183. KANGWAGYE, T. N. 1973. Diurnal and nocturnal biting activity of flies: Diptera in Western Uganda. Bull. Entomol. Res. 63(1): 17-29.
Day and night catches of biting flies were made from bullock, buffalo, and at a light. Species collected were classified according to the time of day at which they bit. Flies caught in early morning included *H. minuta*, *H. spinigera* and *H. t. thirouxi*.

184. ——— 1974. The seasonal incidence of biting flies in Rwenzori National Park and Kigezi Game Reserve, Uganda. Bull. Entomol. Res. 63(4): 535-549.

Biting flies were collected from traps and bait animals. *H. minuta* was the only species to show a unimodal peak. This was attributed to maintenance of soil conditions suitable for the larvae during short rains (September through November). A bimodal peak was shown by *H. spinigera* (minimum numbers in the wet months—September and October).
185. KANO, R. 1954. Nippon no hae. (In Japanese.) DDT Kyokai. pp. 16, 55.
186. ——— 1959. Illustrated insect larvae of Japan. (In Japanese.) Tokyo. p. 692.

Characteristics given for identification of *H. i. exigua*.
187. ——— 1965. Iconographia insectorum Japonicorum colore naturali edita. vol. 3. (In Japanese.) Tokyo, p. 231.
188. ——— S. SHINONAGA, and T. HASEGAWA. 1972. On the specific name of *Haematobia* (Diptera, Muscidae) from Japan. (In Japanese.) Jpn. J. Sanit. Zool. 23(1): 49-56.

H. i. irritans and *H. i. exigua* were distinguishable; the *Haematobia* of Japan believed to be *H. i. irritans*. Characters given to separate *H. i. irritans* from *H. i. exigua*.
189. KARL, O. 1928. 13 Teil. Zweiflugler oder Diptera. III: Muscidae. In Dahl, F. Die Tierwelt Deutschlands und der angrenzenden Meeresteile nach ihren Merkmalen und nach ihrer Lebensweise. Jena, pp. 14-15.
190. KELLOGG, V. L. 1905. American insects. New York, pp. 342-343.
191. KING, H. H. 1911. Report of the entomological section of the Wellcome Tropical Research Laboratories. In Fourth Report of the Wellcome Tropical Research Laboratories at the Gordon Memorial College, Khartoum. Vol. B, General Science (A. Balfour, Ed.) p. 126, plate VI.

Lyperosia (= *Haematobia*) *minuta*, *L. thirourxi* (= *H. t. thirourxi*), and *L. exigua* (= *H. i. exigua*) were collected in Sudan.
192. KINZER, H. G. 1971. Dust-bag and backrubber application of insecticides for control of the horn fly. Vet. Med. Rev. pp. 83-97.
193. ——— and J. M. REEVES. 1974. Dispersal and host location of the horn fly. Environ. Entomol. 3(1): 107-111.

H. i. irritans host location in relation to various environmental factors was investigated. No significant differences were shown between the success of newly emerged and day-old flies in locating hosts or between males and females of either age. In weather conditions, horn fly dispersal and subsequent host location appeared to be random with respect to predominant wind direction. Directional movements, however, were influenced by temperature, wind velocity, and humidity. There was a strong tendency for both parous and nulliparous flies to transfer from one host to another.

194. KIRKWOOD, A. C., and D. W. TARRY. 1973. A survey of some species of flies associated with cattle. *Int. Pest Control* 15(5): 6-10.

H. i. irritans was among various species of flies caught on and near a tethered calf in Surrey, England.

195. KNAPP, F. W. 1972. Prevention of cattle grub infestation in lactating dairy cows by use of daily applications of crotoxyphos. *J. Econ. Entomol.* 65(2): 466-467.

196. ——— 1972. Evaluation of dust bags for horn fly control on cattle. *J. Econ. Entomol.* 65(2): 470-472.

Information on self-dusting stations and the efficacy of several different insecticides was presented. Dusts tested included 3 percent and 5 percent stirofos, 5 percent coumaphos, 3 percent malathion, 5 percent ronnel, 5 percent fenthion, and 3 percent crotoxyphos.

197. KNIPLING, E. F. 1959. The use of DDT in veterinary medicine. *In* S.W. Simmons (ed.). *DDT, the Insecticide Dichlorodiphenyl-trichloroethane and Its Significance*, vol. 2, Basel, pp. 522-528.

198. ——— 1972. Integrated control of livestock insect pests. pp. 379-397. *In* Khan, M. A., and W. O. Haufe. (ed.) *Toxicology, Biodegradation, and Efficacy of Livestock Pesticides. Proceedings of an Advanced Study Institute on Toxicity of Pesticides Used on Livestock Sponsored by the North Atlantic Treaty Organization and organized by the Research Station, Can. Dep. Agric. Lethbridge, Alberta. Amsterdam.* 434 pp.

Discussed possibility of elimination of horn fly in North America.

199. KOE, F-H. 1975. Preliminary observation on the species constitution and the seasonal fluctuations of synanthropic flies in the plain rural region of Honan province China. (In Chinese.) *Acta Entomol. Sin.* 18(1): 71-76.

200. KRAMER, H. 1917. Die musciden der oberlausitz. *Abhandlungen Naturforschenden Gesellschaft, Gorlitz* 28: 289.

201. KRAMER, J. P. 1973. Susceptibility of 16 species of muscoid flies to the microsporidian parasite, *Octospora muscaedomesticae*. J.N.Y. Entomol. Soc. 81(1): 50-53.
H. i. irritans was not susceptible to *O. muscaedomesticae*.
202. KROGERUS, R. 1932. Über die ökologie und verbreitung der arthropen der triebssandgebiete an den Küsten Finnlands. Acta Zool. Fenn. 12:119.
203. KROMBEIN, K. V., and B. D. BURKS. 1967. Hymenoptera of America north of Mexico. Synoptic Catalog. U.S. Dep. Agric., Agric. Monogr. No. 2, 2d Suppl, pp. 254-255.
204. KUNZ, S. E., J. R. CUNNINGHAM, and J. L. ESCHLE. 1973. Horn fly: use of insecticides to disrupt life cycle. J. Econ. Entomol. 66(5): 1239-1240.
Toxicant could reduce or eliminate all infestations of *H. i. irritans* if properly applied to cattle twice in 2 successive weeks. Manure pats were sampled after spraying to test effectiveness of the toxicant.
205. ——— and J. L. ESCHLE. 1971a. Possible use of the sterile male technique for eradication of the horn fly. International Atomic Energy Agency, Food and Agric. Organ., Int. At. Energy Proc. Ser. Sterility Principles for Insect Control or Eradication, pp. 145-156.
206. ——— and J. L. ESCHLE. 1971b. Use of the sterile-male technique to suppress horn flies, *Haematobia irritans* (L.). A pilot study in West Texas. Texas Conference on Insect, Plant Diseases, Weed and Brush Control Proc. 4: 128-130.
207. ——— J. L. ESCHLE, and J. R. CUNNINGHAM. 1975. Methods of estimating sterility in a field population of horn flies (Diptera: Muscidae). J. Med. Entomol. 12(5): 513-517.
Two methods of monitoring the effect of releasing sterile horn flies into a native population of flies were compared. The first method estimated fecundity and hatchability from horn fly eggs collected from individual manure pats. The second method estimated fecundity and hatchability from eggs collected from flies taken from cattle and held in laboratory cages.
208. ——— J. L. ESCHLE, and B. F. HOGAN. 1973. Suppression of a horn fly population with the sterile male technique. Folia Entomol. Mexico. 25-26: 123-124.
209. ——— J. L. ESCHLE, and B. F. HOGAN. 1976. Some bionomical aspects of horn fly populations in West Texas. Southwest. Entomol. 1: 46-48.

210. ——— M. R. GRAHAM, B. F. HOGAN, and J. L. ESCHLE. 1974. Effect of releases of sterile horn flies into a native population of horn flies. *Environ. Entomol.* 3(1): 159-162.
- Low numbers of ^{137}Cs irradiated *H. i. irritans* released into a native population had no effect despite preliminary treatments with insecticide. Larger releases caused a downward trend in reproduction and resulted in about 90 percent control during the latter part of the study.
211. ——— B. F. HOGAN, R. R. BLUME, and J. L. ESCHLE. 1972. Some bionomical aspects of horn fly populations in central Texas. *Environ. Entomol.* 1(5): 565-568.
- Oviposition during October and November was responsible for the spring buildup of flies on cattle during April and May. The average 0.14 flies/pat that emerged from the subsequent overwintering pupae produced an F_1 generation that averaged 1.2 flies/pat. The average monthly rate of production from May to October was 6.6 horn flies/pat when competition from other arthropods was allowed, compared with 66.8/pat when competition was essentially eliminated.
212. LA CHANCE, L. E. 1974. Status of the sterile-insect release method in the world. In *The Sterile-Insect Technique and Its Field Applications*. Proceedings of a panel on the practical use of the sterile-male technique for insect control. FAO/IAEA, Vienna, 1972. pp. 55-62.
213. LAURENCE, B. R. 1953. Some diptera bred from cow dung. *Entomol. Mon. Mag.* 89: 282.
- Lyperosia* (= *Haematobia*) *irritans* reared from cow dung in Harpenden, Herts, England.
214. LEESE, A. S. 1909. Experiments regarding the natural transmission of surra carried out at Mohand in 1908. *J. Trop. Vet. Sci.* 4: 108-132.
- Lyperosia* (= *Haematobia*) and *Haematobia* (= *Haematobosca*) were so scarce that they were believed not to have played an important part in surra transmission.
215. ——— 1912. Biting flies and surra. *J. Trop. Vet. Sci.* 7: 19-34.
- Lyperosia* (= *Haematobia*) *minuta* was believed involved in spread of surra among camels.
216. LEGNER, E. F., and J. H. POORBAUGH. 1972. Biological control of vector and noxious synanthropic flies: a review. *Calif. Vector Views* 19(11): 81-100.
217. LEONARD, M. D. 1928. A list of the insects of New York with a list of the spiders and certain allied groups. *Cornell Univ. Agric. Exp. Stn. Mem.* 101. p. 828.

218. LEVY, R., and H. L. CROMROY. 1973. Concentration of some major and trace elements in forty-one species of adult and immature insects determined by atomic absorption spectroscopy. *Ann. Entomol. Soc. Am.* 66(3): 523-526.
Cu, Fe, Mg, Na, and K content of adult *H. i. irritans* was determined.
219. LLOYD, D. H., and O. O. DIPEOLU. 1974. Seasonal prevalence of flies feeding on cattle in northern Nigeria. *Trop. Anim. Health Prod.* 6(4): 231-236. Abstr. in *Rev. Appl. Entomol.* 63(5): 329-330.
Two species of *Haematobia* were among insects caught in a survey of flies on cattle.
220. LUGGER, O. 1897. Parasites of man and domesticated animals. *Minn. Agric. Exp. Stn. Annu. Rep.* (1896): 188-192.
221. LUTTERMOSER, G. W. 1947. The use of the new insecticides in the control of extoparasites of animals. *Univ. Pa. Bull. (School Vet. Med., Vet. Ext. Q. :107)* 47(28): 68.
222. MA, C. 1959. A preliminary report on the synanthropic flies of Northeastern China. (In Chinese.) *Acta Entomol. Sin.* 9(3): 266.
Lyperosia exigua (= *H. i. exigua*) and *L. titillans* (= *H. t. titillans*) were collected.
223. MACQUEEN, A. 1973. Horn fly breeding, nitrogen loss, and nutrient immobilization associated with cattle dung in the southern interior of British Columbia. Simon Fraser Univ. Ph.D. Dissertation. 213 pp.
224. ——— and B. P. BEIRNE. 1974. Insects and mites associated with fresh cattle dung in the southern interior of British Columbia. *J. Entomol. Soc. Brit. Columbia* 71: 5-9.
H. i. irritans was among the 67 insect species found to be associated with dung in B.C.
225. ——— and B. P. BEIRNE. 1975a. Dung burial activity and fly control potential of *Onthophagus nuchicornis* (Coleoptera: Scarabaeinae) in British Columbia. *Can. Entomol.* 107(11): 1215-1220.
Beetle activity in the field had little effect on the horn fly as the beetles ceased burial activity before midsummer when horn flies reached their greatest numbers.
226. ——— and B. P. BEIRNE. 1975b. Influence of other insects on production of horn fly, *Haematobia irritans* (Diptera: Muscidae) from cattle dung in south-central British Columbia. *Can. Entomol.* 107(12): 1255-1264.

Horn flies did not show a definite diurnal rhythm in oviposition. There was a single peak adult abundance during the summer. Predators *Philonthus cruentatus* Gmelin, *Sphaeridium scarabaeoides* (L.), and *S. lunatum* (F.) were chiefly responsible for horn fly suppression in B.C. Horn fly parasites and dung burying beetles were scarce.

227. MAGYAR, K., and G. MAKARA. 1974. On the possibilities of preventing the damage caused by flies in cattle. (In Hungarian.) Magy. Allatorv. Lapja 29(7): 447-448, 453-456.

228. MALVIYA, H. C. 1972. Stephanofilarial infection in cattle and buffalo in Anadaman Islands. Indian J. Helminthol. 24(2): 68-71.

Of the 1,002 flies collected from lesions on cattle and buffalo, 64.6 percent were *H. i. exigua*.

229. MAYER, M. S., D. L. SILHACEK, D. A. CARLSON, and J. D. JAMES. 1972. Attraction of the male house fly to cuticular hydrocarbons and feces of several other dipteran species. Experientia (Basel) 28(9): 1111-1112.

Musca domestica was attracted to *H. i. irritans* cuticular hydrocarbons.

230. MAYER, R. T., and A. C. BRIDGES. 1975. Some effects of ionizing radiation on the lipid and glycogen content of adult horn flies, *Haematobia irritans*. Insect Biochem. 5(4): 387-398.

Measurements were made of the free sugar, glycogen, and lipid levels in flies treated with a 2.0 krad dose of ionizing radiation. Glycogen levels were depressed in males but remained unchanged in females. Triglyceride levels in both sexes were 50 percent lower than the control group, but there was no effect on the distribution of fatty acids among the various lipid classes.

231. ——— J. COOPER, F. M. FARR, and R. H. SINGER. 1975. Some effects of ionizing radiation on adult horn flies, *Haematobia irritans*. Insect Biochem. 5(1): 35-42.

Adult flies were irradiated from a ¹³⁷Cs source and analyzed for physiological (free) amino acids and for total (hydrolysates) amino acids. Assays showed that irradiated flies had increased levels of asparagine—glutamine and creatinine—but neither ammonia nor total amino acids were affected. Several compounds previously unreported in insects were found to occur in horn flies.

232. ——— F. M. FARR, and J. COOPER. 1975. The occurrence of creatinine and creatine phosphate in adult horn flies. J. Ga. Entomol. Soc. 10(1): 61-64.

Extracts of whole adult horn flies contained 11.0 μ -mole creatinine and 0.5-1 μ -mole creatine phosphate per gram tissue.

233. MAXWELL, F. G. 1975. Department of entomology annual report for 1973-1974. Miss. Agric. Forest Exp. Stn., Res. Highlights 38(1): 8.
234. MAXWELL-LEFROY, H. 1907. A preliminary account of the biting flies of India. Agric. Res. Inst., Pusa. Bull. No. 7: 31, 35-38.
235. ——— and F. M. HOWLETT. 1909. Indian insect life. Calcutta, pp. 643-647.
236. McKEOWN, K. C. 1944. Australian insects. Sydney, pp. 233-234.
237. MEIGEN, J. W. 1838. Systematische beschreibung der bekannten Europaischen zweiflugeligen insekten, vol. 7, p. 171.
238. MELVILLE, R. V. 1974. *Siphona* Meigen, 1803, and *Haematobia* Lepeletier and Serville, 1828 (Insecta: Diptera); designations of type-species under the plenary powers. Bull. Zool. Nomencl. 30(3/4): 157-158.

Musca geniculata DeGeer, 1776 was designated type-species for *Siphona* Meigen, 1803. Designation by Westwood (1840) of *Conops irritans* Linnaeus as type-species of *Haematobia* Lepeletier and Serville, 1828 was confirmed.
239. MENG, C. H., and G. F. WINFIELD. 1950. Studies on the control of fecalborne diseases in North China, XVI. An approach to the quantitative study of the house-frequenting fly population. D. The breeding habits of the common North China flies. Philipp. J. Sci. 79: 165-192.
240. MERRITT, R. W. 1974. The species diversity and abundance of insects inhabiting cattle droppings and their role in the degradation of dung in four different pasture and rangeland ecosystems in the Sierra Nevada foothills of California. Ph.D. Dissertation, Univ. Calif., Berkeley. 273 pp.
241. ——— 1975. Cattle dropping degradation activities by the scarab, *Aphodius fimetarius* (L.), in the Sierra foothills of California. North Cent. Branch Entomol. Soc. Am. Proc. 30: 97.

Presence of *A. fimetarius* adults in cow pats reduced the number of horn flies emerging from the droppings.
242. ——— and J. H. POORBAUGH. 1975. A laboratory collection and rearing container for insects emerging from cattle droppings. Calif. Vector Views 22(6): 43-46.
243. MESSENGER, P. S. 1959. Bioclimatic studies with insects. Annu. Rev. Entomol. 4: 192.
244. MILLER, J. A., J. L. ESCHLE, and I. L. BERRY. 1969. Patterns of flight activity in livestock insects. 1. Preliminary testing of a system for recording flight activity of the stable fly. Ann. Entomol. Soc. Am. 62(5): 1046-1050.

245. MILLER, R. W., and L. G. PICKENS. 1973. Evaluation of stirofos as a feed additive larvacide for fly control on dairy farms. (Abstr.) J. Dairy Sci. 56(5): 669.
H. i. irritans were controlled with stirofos fed to cattle at rate of 1.5 mg/kg.
246. ——— and L. G. PICKENS. 1975a. Feed additives for control of flies on dairy farms. J. Med. Entomol. 12(1): 141-142.
1.5 mg stirofos per kg body wt controlled horn flies.
247. ——— and L. G. PICKENS. 1975b. Evaluation of methoprene formulations for fly control. J. Econ. Entomol. 68(6): 810-812.
Neither encapsulation nor a slow release plastic formulation improved the effectiveness of methoprene tested as a feed additive for fly control. Salt blocks that contained 0.02 percent methoprene appeared to reduce horn flies on pastured dairy heifers.
248. ——— L. G. PICKENS, N. O. MORGAN, R. W. THIMIJJAN, and R. L. WILSON. 1973. Effect of stable flies on feed intake and milk production of dairy cows. J. Econ. Entomol. 66(3): 711-713.
249. ——— and F. C. UEBEL. 1974. Juvenile hormone mimics as feed additives for control of the face fly and house fly. J. Econ. Entomol. 67(1): 69-70.
250. MITTER, J. L. 1916. The life history of *Haematobia sanguisuga* Austen. Indian J. Med. Res. 3: 530-537, 538-540.
251. MORGAN, C. E., and G. D. THOMAS. 1974. Annotated bibliography of the horn fly, *Haematobia irritans* (L.), including references on the buffalo fly, *H. exigua* (de Meijere), and other species belonging to the genus *Haematobia*. U.S. Dep. Agric. Misc. Publ. 1278. 134 pp.
252. NAGASAWA, S. 1959. Biological assay of insecticide residues. Annu. Rev. Entomol. 4: 334.
253. NEAVE, N. A. 1912. Notes on the blood-sucking insects of eastern tropical Africa. Bull. Entomol. Res. 3(3): 310, 317-318, 320, 322.
Lyperosia (= *Haematobia*) was found feeding on freshly killed animals.
254. NEILSON, C. L. 1955. Handbook of the main economic insects of British Columbia. Part 2. Livestock insects. Unpubl. Mimeo. B.C. Dep. Agric., 40 pp.
255. NEVEU-LEMAIRE, M. 1938. Traite D'Entomologie medicale et veterinaire. Paris, pp. 943-945.
256. NEVILL, E. M. 1975. Preliminary report on the transmission of *Parafilaria bovicola* in South Africa. Onderstepoort J. Vet. Res. 42(1): 41-48.

257. NEWTON, W. H., P. J. HAMMAN, and M. A. PRICE. 1973. Suggestions for controlling external parasites of livestock and poultry. Tex. A & M Univ. Agric. Ext. Serv. MP-691, 40 pp.
258. NUORTEVA, M. 1945. Observations on the life of *Oxybelus uniglumis* L. (Hym., Sphegidae). (In Finnish.) Ann. Entomol. Fenn. 11(4): 216.
H. (= *Haematobosca*) *stimulans* was preyed upon by *O. uniglumis*.
259. OLDROYD, H. 1964. The natural history of flies. London, 324 pp.
260. OSMUN, J. V. 1964. Insects and other arthropods of economic importance in Indiana in 1963. Proc. Indiana Acad. Sci. 72: 146-153.
261. PAINTER, R. H. 1953. Horn-fly control on beef cattle in the prairie provinces. Can. Dep. Agric., Sci. Serv.-Div. Entomol. 6 pp.
262. PAL, R. and L. E. LACHANCE. 1974. The operational feasibility of genetic methods for control of insects of medical and veterinary importance. Annu. Rev. Entomol. 19: 269-291.
263. PAPP, L. 1971. Ecological and production biological data on the significance of flies breeding in cattle droppings. Acta Zool. Acad. Sci. Hung. 17(1-2): 91-105.
264. PATE, F. M., and M. J. JANES. 1975. Use of Trolene-40 for control of horn flies and cattle grubs. Mimeo. Rep. Fla. Everglades Exp. Stn. (Belle Glade) 3: 7 pp.
265. PATNAIK, B., and V. KUMAR. 1972. A research note on the occurrence of juveniles of *Stephanofilaria*, the causative parasite of ear sore in buffaloes, in *Musca autumnalis*. Indian J. Anim. Sci. 42(5): 351-352.
266. PATTON, W. S. 1925. Diptera of medical and veterinary importance, I. Types of older authors in continental museums. Philipp. J. Sci. 27(2): 177-200.
 Synonymy of *Haematobia* was discussed.
267. ——— and E. MACGILL. 1925. Comparative studies of the antennae of some of the higher Diptera. Indian J. Med. Res. 13: 284-285.
268. PECK, O. 1963. A catalogue of the Nearctic Chalcidoidea (Insecta: Hymenoptera). Can. Entomol. Suppl. 30. 1,092 pp.
 Included various parasitic chalcids of *H. i. irritans*.
269. ——— 1974. Chalcidoid parasites of the horn fly, *Haematobia irritans* (Diptera: Muscidae), in Alberta and elsewhere in Canada. Can. Entomol. 106(5): 473-477.

- Horn fly was parasitized by *Muscidifurax raptor*, *M. zaraptor*, *Eupteromalus viridescens*, *Spalangia drosophilae*, *S. subpunctata*, and *S. e. erythromera*. *Spalangia* spp. was credited with parasitizing up to 40 percent of horn fly pupae in traps, and *Muscidifurax* spp. up to 30 percent.
270. PERIS, S. V. 1951. Los Stomoxydinae de la Peninsula Iberica (Dipt., Muscidae). Rev. Acad. Cienc. Exactas Fis-Quim. Nat. Zaragoza 6: 41-49.
Discussed the classification of Spanish stomoxiine flies.
 271. ——— 1960. Notas Dipterologicas. Graellsia 18: 59-63.
Discussed stomoxiine flies found in Spain.
 272. PETERSON, A. 1960. Larvae of insects. Part II. Ann Arbor, pp. D27, 260, 316.
Identification of horn fly larvae is presented.
 273. PICKENS, L. G., and R. W. MILLER. 1975. Growth-regulating chemicals tested against nontarget insect fauna in bovine fecal pats. Environ. Entomol. 4(1): 46.
 274. PONT, A. C. 1970. The type material of Oriental and Australasian Muscidae (Diptera) in the Zoological Museum, Amsterdam. Beaufortia 18(231): 86.
H. i. exigua lectotype, paralectotype, and seven syntypes were present in the collection.
 275. ——— 1973a. Muscidae (house-flies, stable-flies etc.). In Smith, G. V. (ed.). Insects and Other Arthropods of Medical Importance. Brit. Museum (Nat. Hist.) London, pp. 251-269, 489.
 276. ——— 1973b. Studies on Australian Muscidae (Diptera). IV. A revision of the subfamilies Muscinae and Stomoxyinae. Aust. J. Zool. Suppl. Ser. 21: 129-296.
H. i. exigua was regarded as a subspecies of *H. irritans* in accordance with Zumpt (1973).
 277. POPHAM, W. L., and D. G. HALL. 1958. Insect eradication programs. Annu. Rev. Entomol. 3: 336.
 278. POVOLNY, D. 1959. Gesichtspunkte der klassifikation von synanthropen fliegen. Z. Angew. Zool. 46: 324-328.
 279. RACK, G. 1975. Two new species of the genus *Pediculaster* from Australian Diptera. (In German.) Acarologia (Paris) 16(3): 500-505.
P. domrowi sp. n. and *P. morelliae* sp. n. were found attached at the halteres, thorax, and abdomen of *H. i. exigua* and *Morellia hortensia*.
 280. RAILLIET, A. 1886. Elements de Zoologie Medical et Agricole. Paris, p. 548.

281. RANSOM, B. H. 1908. The animal parasites of cattle. U.S. Dep. Agric., Bureau of Animal Industry. Special Rep. on Diseases of Cattle. pp. 496-497.
282. REID, E. T. M. 1956. Notes on the distribution of Stomoxydinae (Dipt., Muscidae) in the Southern Sudan. Entomol. Mon. Mag. 92: 343-346.
283. REINHARDT, R. 1975. Shoo fly. Zoecon Corp. (Palo Alto, Calif.) Annu. Rep. for year ended December 31, 1974, pp. 5-10.
Discussed possible control of horn fly with growth regulator methoprene.
284. RIEDEL, M. P. 1919. Beitrag zur kenntnis der dipterenfauna des Niederrheins. Entomol. Zeitschr. Frankfurt, 33: 39.
285. RINGDAHL, O. 1921. De skanska stranddynernas insektfauna. Entomol. Tidskr. 41: 39.
286. ——— 1926a. Neue nordische musciden nebst berichtigung und namensanderungen. Entomol. Tidskr. 47: 101.
287. ——— 1926b. Musciden und Anthomyiiden von estlandischen hochmooren und nachtrag zu den Tachiniden. In Dampf, A. Zur Kenntnis der estlandischen Hochmoorfauna. Sitzungberichte der Naturforscher-Gesellschaft bei der Universitat Tartu 33: 86.
288. ——— 1928. Beitrage zur kenntnis der Anthomyidenfauna des nordlichen Norwegens. Tromso Museums Arshefter 49(3): 10.
289. ——— 1945. Oversikt av smaslaktena inom familjen Muscidae. Entomol. Tidskr. 66: 95-106.
290. ——— 1951. Flugor fran fjalltrakterna. Opusc. Entomol. 16: 171.
291. ——— 1952. Catalogus insectorum Sueciae. Diptera, Cyclorapha: Muscaria, Schizometopa. Opusc. Entomol. 17: 150.
292. ——— 1959. Flugor pa strandangar i nordvastra skane. Entomol. Tidskr. 80: 44.
293. ——— 1960. Flugfaunan pa kullaberg och hallands vaderö. Kullabergs Natur 2: 24.
294. RIVOSCECHI, L. 1953a. Contributo alla conoscenza delle Stomoxydinae Italiane (Diptera: Muscidae). Riv. Parassitol. 14: 29-36.
295. ——— 1953b. I Muscidae pungenti (Diptera, Stomoxydinae) Italiani. Rend. Inst. Super. Sanita 16: 286-332.
Morphological and ethological aspects of five species of Stomoxyine flies were studied.

296. ——— and E. STELLA. 1974. Bloodsucking arthropods from nature reserve areas. Part 1. The Circeo National Park, Italy. (In Italian.) *Parassitologia*. Rome, 15(3): 281-296.
297. ROARK, R. C., and N. E. McINDOO. 1944. A digest of the literature on DDT through April 30, 1944. U.S. Dep. Agric., Entomol. and Plant Quarantine. E-631. pp. 43-44.
298. ROBERTS, J. E. 1969. Horn fly control. *Arkansas Cattle Bus.* 5(7): 8.
299. ROBERTS, R. H. 1972a. Horn fly control with insecticides applied with photo-actuated sprayers. *Miss. Agric. Exp. Stn. Inform. Sheet 1181*, 2 pp.
300. ——— 1972b. Sprayer control of horn flies. *Miss. Farm. Res.* 35(3): 4-6.
301. ——— and R. A. HOFFMAN. 1970. Horn flies on cattle controlled with several insecticides applied with a power spray. *Miss. Agric. Exp. Stn. Inform. Sheet 1112*, 2 pp.
302. ——— and W. A. PUND. 1974. Control of biting flies on beef steers: effect on performance in pasture and feedlot. *J. Econ. Entomol.* 67(2): 232-234.

Steers from insecticide treated groups averaged higher daily gains and required less feed per pound of gain than steers from the control groups, but the differences were not significant.
303. RODRIGUEZ, J. G., J. SINGH, and B. TAYLOR. 1970. Manure mites and their role in fly control. *J. Med. Entomol.* 7: 335-341.
304. RONALD, N. C., and C. W. WINGO. 1973. Cost and effectiveness of horn fly and face fly control programs in semi-isolated range herds in central Missouri. *J. Econ. Entomol.* 66(3): 693-696.

Standard control programs and a ronnel-supplement program were used under large herd-large pasture conditions. Only sprays produced satisfactory control of *H. i. irritans* without supplementary control measures.
305. ROSELLE, R. E. 1956. Control livestock insects. *Univ. Nebr. Ext. Circ.* 56-1575, 2 pp.
306. ——— and J. B. CAMPBELL. 1970. Horn flies on range cattle. *Nebr. Cattleman* 26(9): 36.
307. ROUBAUD, E. 1906. Stomoxides nouveaux du Senegal. *Cent. Rech. Seances Soc. Biol.* 60: 895-897.
308. ——— 1907. Stomoxides nouveaux du Congo. *Ann. Inst. Pasteur (Paris)* 21: 666-669.
309. RUZIMURADOV, A. 1972. Protective action of Co-Ral and Dibrom against pasture flies. (In Russian.) *Tr. Uzb. Nauchno-Issled. Inst. Vet.* 20: 135-136.

310. ——— 1974. Flies, parasites of skin integument of domestic animals. (In Russian.) *Parazitologiya* (Leningr.) 8(5): 447-448.
311. ——— and T. PIRNAZAROV. 1972. Flies of farm stock in the Kashkadar' Insoki regions. (In Russian.) *Tr. Uzb. Nauchno-Issled. Inst. Vet.* 20: 137.
312. SABROSKY, C. W. 1971. The type-species of *Siphona* Meigen, 1803, and *Haematobia* Lepeletier and Serville, 1828 (Insecta: Diptera) *Z.N. (S.)* 195. *Bull. Zool. Nomencl.* 27(5-6): 234-237.
313. SANCHEZ, S. A. 1973. Dung beetles. *Cattleman* 59(10): 76-77.
314. SAVIO, T., L. HAHN, H. D. JOHNSON, and G. D. THOMAS. 1972. Effect of horn flies on 3-methoxy-4-hydroxy mandelic acid excretion in cattle. *J. Dairy Sci.* 55(5): 700. (Abstract only).
Increased excretion of vanilmandelic acid believed to result from increase in standing time and nervous activity associated with the physical disturbance caused by the biting of the flies.
315. SAVIO, T. J., H. D. JOHNSON, L. HAHN, and G. D. THOMAS. 1976. Effect of horn flies on vanilmandelic acid excretion of dairy cattle. *J. Dairy Sci.* 59(2): 318-320.
Increased vanilmandelic acid excretion in urine reflected an increase in standing time and nervous activity associated with the physical disturbance caused by the biting activity of the flies. Study was conducted on six lactating Holstein cows exposed to approximately 500 horn flies per cow per day for 4 weeks.
316. SCHLINKE, J. C., and J. S. PALMER. 1973. Combined effects of phenothiazine and organophosphate insecticides in cattle. *J. Am. Vet. Med. Assoc.* 163(7. pt. 1): 756-758.
At recommended dosages, no potentiation of the toxic effect of any of the insecticides by phenothiazine existed.
317. SCHMIDT, C. D. 1972. Classification of the physiological development of laboratory-reared female horn flies, *Haematobia irritans*. *Ann. Entomol. Soc. Am.* 65(3): 695-701.
The following characteristics were used to classify physiological age or development: presence of sperm in spermathecae, presence of follicular relics, pattern of tracheoles on ovarioles, contents of malpighian tubes, appearance of fat bodies, and numbers of nonfunctional ovarioles.
318. ——— and R. R. BLUME. 1973. Laboratory-reared horn flies: relationships between width of head and weight of pupa in both sexes and between these measurements and number of ovarioles in females. *Ann. Entomol. Soc. Am.* 66(6): 1307-1308.

Head width at the widest point was significantly correlated with the weight of 6-day-old pupae. The average number of ovarioles/female was correlated with the average weight of the pupa and the average width of the head.

319. ——— J. L. ESCHLE, S. E. KUNZ, and J. M. DREISS. 1974. Sterile-male technique for suppression of the horn fly: effects of irradiation, handling, and release on laboratory-reared flies. *Environ. Entomol.* 3(2): 287-289.
Emergence of adults was reduced slightly by irradiation and significantly by shipment and release. Refrigeration to delay emergence of adults reduced percentage emergence.
320. ——— C. R. WARD, and J. L. ESCHLE. 1973. Rearing and biology of the horn fly in the laboratory: effects of density on survival and fecundity of adults. *Environ. Entomol.* 2(2): 223-224.
When 100 adult flies were placed in cages of 7 sizes or various numbers of flies were placed in 237-cm³ cages, the hatchability of the eggs obtained from the more densely populated cages was always higher. Survival of flies was higher when the cages were less densely populated. Densities of caged flies ranging from 0.6 to 9.5 cm³ of space/fly were recommended for rearing laboratory flies.
321. SCHMIDT, F. M. 1971. Ranchers adopt aerial spraying for flies. Serving Farm, Ranch, Home. *Univ. Nebr. Coll. Agric. Home Econ. Q.* 18(2): 15-16.
322. SCHNABL, J. 1911. Dipterologische sammelreise nach Korsika. *Dtsch. Entmol. Z.* p. 97.
323. SCHUMANN, H. 1961. Die eier von fliegen. *Mikrokosmos* 50: 297-300.
324. ——— 1963. Zur larvalsystematik der Muscinae nebst beschreibung einiger musciden und anthomyidenlarven. *Dtsch. Entomol. Z.* 10: 134-163.
Discussed larval stages of various muscine flies including *H. i. irritans* and *H. t. titillans*. Includes key for separating third stage muscid larvae and illustrations of cephalopharyngeal skeletons and spiracular plates for various species of muscids.
325. SCHWARTZ, B., R. E. REBRASSIER, J. E. ACKERT, and others. 1947. Report of the committee on parasitic diseases. *U.S. Livestock Sanit. Assoc. Rep.* 50: 98.
326. SCHWARZ, M., R. W. MILLER, J. E. WRIGHT, and others. 1974. Compounds related to juvenile hormone. Exceptional activity of arylterpenoid compounds in four species of flies. *J. Econ. Entomol.* 67(5): 598-601.

The six compounds tested were highly active against all four species of flies including the horn fly.

327. SCOTT, H. G., and K. S. LITTIG. 1962. Flies of public health importance and their control. Insect Control Series: Part 5. U.S. Dep. Health Educ. Welfare, Public Health Serv. Publ. No. 772, 40 pp.
328. SEGUY, E. 1924. Les insectes parasites de l'homme et des animaux domestiques. Encyclopedie Pratique du Naturaliste. Paris, pp. 276-280.
329. ———. 1933. Contributions a l'etude de la faune du Mozambique. Voyage de M. P. Lesne (1928-1929). Memorias e estudos do Museu Zoologico da Universidade de Coimbra. Serie 1. No. 67. p. 54.
330. ———. 1934. Dipteres (Brachyceres) (Muscidae Acalypterae et Scatophagidae). Faune Fr. 28: 25.
331. ———. 1937. Diptera. fam. Muscidae. fasc. 205. In Wytsman, P. (ed.) Genera Insectorum. Bruxelles. 604 pp.
Discussed synonymy, types, and geographical distribution of *Haematobia* and other stomoxine flies.
332. ———. 1951. Ordre des Dipteres. In Traite de Zoologie. Anatomie, Systematique, Biologie. Tome 10. Insectes Superieures et Hemipteroides (Premier Fascicule). Paris, pp. 721-722.
333. ———. 1957. Anthophilie des dipteres haematophages. Rev. Fr. Entomol. 24: 390.
334. SEN, S. K., and T. B. FLETCHER. 1962. Veterinary entomology and acarology for India. New Delhi, 668 pp.
335. SENIOR-WHITE, R. 1922. Notes on Indian Diptera. Mem. Dep. Agric. India. Entomol. Ser. 7(9): 99.
336. SHINONAGA, S., and R. KANO. 1971. Fauna Japonica, Muscidae I (Insecta: Diptera). Tokyo, pp. 87-90.
Discussed taxonomy and geographical distribution of *H. i irritans* and *H. i. exigua* in Japan.
337. SHOL', V. A. 1969. Description of the causative organism of setariosis in marals. (In Russian.) Izv. Akad. Nauk Kaz. SSR Ser. Biol. 6:45-50.
338. ——— and N. I. DROBISHCHENKO. 1971. Development of the causative agent of setariosis of marals in the body of the blood-sucking fly, *Haematobia stimulans*. (In Russian.) Dokl. Akad. Nauk SSSR Ser. Biol. 199 No. 2, pp. 503-504.
339. SHULL, W. E., and R. A. FISHER. 1940. Recommendations for insect control. Idaho Ext. Bull. No. 129, p. 31.

340. SIMPSON, J. J. 1911. Entomological research in British West Africa. I. Gambia. Bull. Entomol. Res. 2: 187-239.
341. SINGH, A., and R. C. CHHABRA. 1973. Incidence of arthropod pests of domesticated animals and birds. Indian J. Anim. Sci. 43(5): 393-397.
342. SMART, J. 1948. A handbook for the identification of insects of medical importance. Brit. Museum (Nat. Hist.) London, pp. 61-62.
343. SMITH, J. B. 1900. Insects of New Jersey. Suppl. 27th Annu. Rep. N.J. State Board of Agric., pp. 559, 678-679.
Spalangia haematobiae was listed as a parasite of the horn fly.
344. SNYDER, F. M. 1965. Insects of Micronesia, Diptera: Muscidae. Insects Micronesia 13(6): 311-313.
H. i. exigua was found on islands of S. Mariana, Palau, Yap, and Ponape.
345. SPACKMAN, E. W., and J. E. LLOYD. 1975. Control insect pests of beef cattle, 1975. Wyo. Agri. Exp. Stn. Bull. 544 R. 20 pp. (Rev. Annu.)
346. STAAL, G. B. 1975. Insect growth regulators with juvenile hormone activity. Annu. Rev. Entomol. 20: 427.
347. STANDFAST, H. A., and A. L. DYCE. 1972. Arthropods biting cattle during an epizootic of ephemeral fever in 1968. Aust. Vet. J. 48(3): 77-80.
348. STEELMAN, C. D. 1976. Effects of external and internal arthropod parasites on domestic livestock production. Annu. Rev. Entomol. 21: 155-178.
349. ——— T. W. WHITE, and P. E. SCHILLING. 1972. Effects of mosquitoes on the average daily gain of feedlot steers in Southern Louisiana. J. Econ. Entomol. 65(2): 462-466.
350. STOFFOLANO, J. G. 1970. Nematodes associated with the genus *Musca* (Diptera: Muscidae). Bull. Entomol. Soc. Am. 16(4): 194-203.
351. STORK, M. N. 1936. A contribution to the knowledge of the puparia of Anthomyidae. Tijdschr. Entomol. 79: 98, 100, 113-115.
The puparia of *H.* (= *Haematobosca*) *stimulans* and *Lyperosia* (= *Haematobia*) *irritans* were among the 35 species of puparia discussed.
352. STRICKLAND, E. H. 1952. Insects of the year in Northern Alberta, 1951. Can. Insect Pest Rev. 30(1): 24.


353. STROBL, P. G. 1899. Spanische dipteren. Wiener Entomol. Zeitung 18: 218.
Stomoxys (= *Haematobia*) *irritans* was found near Madrid.
354. ——— 1900. Dipterenfauna von bosien, hercegovina und dal-matien. Wissenschaftliche Mittheilungen aus Bosnien und Hercegovina. vol. 7, pt. 3, p. 607.
355. SURCOUF, J., and R. GONZALES-RINCONES. 1911. Essai sur les Dipteres vulnerants du Venezuela, I, Nematoceres; II, Brachy-ceres. Paris.
356. SURFACE, H. A. 1913. Pests of domestic animals. Bi-monthly Zool. Bull., Div. of Zool., Pa. Dep. Agric. 3(1): 14-15.
357. SYCHEVSKAYA, V. I. 1956. Flies of medical importance in Kara-Kalpakia. (In Russian.) Entomol. Obozr. 35: 352.
358. ——— 1957. Flies of medical importance of N. Kirghisia. (In Russian.) Entomol. Obozr. 36: 110, 112-114.
359. ——— 1963. Some Chalcidoidea reared from puparia of synan-thropic flies in Uzbekistan. Zool. Zh. 42: 858-864.
Haematobia was mentioned as being the host for several of the seven species of parasites used in the study.
360. SZTANKAY-GULYAS, M. 1953. Muscidae in Hungary and sur-rounding areas. Acta Vet. Acad. Sci. Hung. 3: 245-286.
 Included a discussion of several *Haematobia* species. Presented morphological descriptions, keys, geographical distribution, and a short characterization for each species.
361. TIENSUC, L. 1935. Die bisher aus Finnland bekannten musciden. Acta Soc. Fauna Flora Fenn. 58(4): 53.
362. ——— 1948. On the phenology of flower visiting by Diptera in southern Finland. (In Finnish.) Ann. Entomol. Fenn. 14(Suppl.): 216.
363. TILLYARD, R. J. 1926. The insects of Australia and New Zea-land. Sydney, p. 375.
364. TIMON-DAVID, J. 1937. Recherches sur le peuplement des hautes montagnes. Dipteres de la vallee de Chamonix et du massif du Mont Blanc. Ann. Fac. Sci. Mars. 3: 11.
H. (= *Haematobosca*) *stimulans* (Meigen) was among the in-sects collected.
365. ——— 1950. Dipteres des Pyrenees. Bull. Soc. Hist. Nat. Toulouse 85: 16.
366. THOMAS, G. D., I. L. BERRY, and C. E. MORGAN. 1974. Field developmental time of non-diapausing horn flies in Missouri. Environ. Entomol. 3(1): 151-155.

An equation was formulated to predict field developmental time of *H. i. irritans*: $1/T = -0.04769 + 0.005168 \theta$; T = time required for development (in days) and θ = mean temperature ($^{\circ}\text{C}$) during development.

367. THOMPSON, W. R. 1955. A catalogue of the parasites and predators of insect pests. Sec. 2, Pt. 3. Hosts of the Hymenoptera (Calliceratid to Evaniid). Commonw. Agric. Bur. Publ. p. 213.
 368. ——— 1958. A catalogue of the parasites and predators of insect pests. Sec. 2, Pt. 5. Hosts of the Hymenoptera (Miscogasteridae to Trigonalidae) Lepidoptera and Strepsiptera. Commonw. Agric. Bur. Publ. pp. 620-622.
 369. THOMSEN, E. G., and M. H. DONER. 1938. Fly control for livestock. Soap 14(7): 101-103.
 370. TORGERSEN, T. R. 1974. Host index to the second supplement of the synoptic catalog for the Hymenoptera of America north of Mexico. Bull. Entomol. Soc. Am. 20(4): 288.
 371. TOWNSEND, C. H. T. 1932. Note on *Haematobia* and *Lyperosia*. Rev. Entomol. (Rio de J.) 2: 241.
 372. ——— 1937. Genera of Glossinidae to Muscidae. Manual of myiology. V. pp. 23-26.
 373. TWINN, C. R., and F. A. HERMAN. 1928. A cheap and effective fly spray. 58th (1927) Entomol. Soc. Ont., Annu. Rep. p. 44.
 374. TYNES, J. S., E. C. BURNS, and C. E. SEVERENCE. 1970. Dust bags for horn fly control. La. State Univ. Ext. Publ. 1613, 6 pp.
 375. TZE-TEH, FAN. 1957. Notes on third stage larvae of synanthropic flies in Shanghai district. (In Chinese.) Acta Entomol. Sin. 7(4): 405-422.
- Key presented for 31 species including *Lyperosia* (= *Haematobia*) *i. exigua*.
376. UNITED STATES DEPARTMENT OF AGRICULTURE. 1952. Some important insects. U. S. Dep. Agric., Yearb. 1952. Plate XXV.
 377. ——— 1969. Horn flies on cattle, how to control them. Leaflet No. 388, 8 pp. (Rev. Annu.)
 378. ——— 1971. Horn fly, *Haematobia irritans*. U.S. Dep. Agric., Coop. Econ. Insect Rep. 21: 26, 83, 127, 190, 212, 235, 239, 264, 296, 312, 326, 346, 362, 380, 399, 417, 436, 456, 476, 496, 515, 533, 552, 555, 570, 591, 606, 623, 634, 662, 685, 698, 711, 723, 742, 754.
 379. ——— 1972. Horn fly, *Haematobia irritans*. U.S. Dep. Agric., Coop. Econ. Insect Rep. 22: 147, 167, 174, 175, 193, 215, 230, 244, 256, 271, 281, 308, 321, 336, 359, 389, 410, 428, 442, 464, 485, 504, 528, 545, 564, 581, 605, 623, 641, 663, 683, 697, 713, 725, 740, 753, 764.

380. ——— 1973. Horn fly, *Haematobia irritans*. U.S. Dep. Agric., Coop. Econ. Insect Rep. 23: 5, 129, 148, 155, 158, 169, 184, 204, 213, 238, 257, 272, 289, 306, 324, 345, 365, 386, 387, 405, 428, 448, 470, 491, 516, 539, 562, 603, 621-622, 643, 658, 673, 689, 704, 715, 728, 742, 743, 758, 779.
381. ——— 1974. Horn fly, *Haematobia irritans*. U.S. Dep. Agric., Coop. Econ. Insect Rep. 24: 8, 48, 73, 94, 104, 132, 169, 201, 228-229, 250, 254, 258-259, 275-276, 303, 327, 346, 370, 373, 391, 413, 434, 463-464, 487, 523, 524, 551, 581, 609, 633-634, 657, 680, 701, 746, 759, 774, 776, 788, 802, 813, 828, 842, 857, 872.
382. ——— 1975. Horn fly, *Haematobia irritans*. U.S. Dep. Agric., Coop. Econ. Insect Rep. 25: 134, 173, 209, 244, 268, 272, 276-277, 294, 320, 334, 352, 373, 395, 417, 448, 469, 490-491, 512, 539, 560, 580, 601, 624, 647, 669, 689, 709, 727, 746, 749, 761, 772, 784, 795, 810, 821, 833, 846, 874, 899, 901.
383. URSU, A. 1968. New contributions to the knowledge of the synanthropic Diptera fauna of the Socialist Republic of Romania. (In Romanian.) pp. 409-410. In Cadariu, G. H., and B. Barhad. Activitatea Stintifica a Institutului de Igiena, Bucharesti 1927-1967. Editura Med. Icala, Bucharest, 457 pp.
384. VIERECK, H. L. 1916. The Hymenoptera, or wasp-like insects, of Connecticut. Part III. Guide to the insects of Connecticut. Conn. State Geol. Nat. Hist. Surv. Bull. 22: 485-486.
Spalangia haematobiae was reared from larva of the horn fly.
385. WALKER, F. 1849. List of the specimens of dipterous insects in the collection of the British Museum. Part III. London. p. 681.
386. WASHBURN, F. L. 1919. The hymenoptera of Minnesota. 17th Annu. Rep. (1918) State Entomol. Minn. p. 187.
Listed several parasites of *H. i. irritans*.
387. WATERHOUSE, D. F. 1974. The biological control of dung. Sci. Am. 230(4): 100-109.
388. WATTS, K., and R. L. COMBS. 1975. New host record for *Aleochara bipustulata*. J. Econ. Entomol. 68(4): 564.
389. WEAVER, G. S., J. J. O'CONNELL, and G. I. GILBERTSON. 1946. Spraying cattle insects. South Dakota Ext. Leaflet No. 102. pp. 5-6.
390. WEINER, T. J., and E. J. HANSSENS. 1975. Species and numbers of bloodsucking flies feeding on hogs and other animals in southern New Jersey. J.N.Y. Entomol. Soc. 83(3): 198-202.
H. i. irritans was observed feeding on horses.
391. WEST, L. S. 1951. The housefly: its natural history, medical importance and control. Ithaca, pp. 195, 198, 261.

392. ——— and O. B. PETERS. 1973. Annotated bibliography of *Musca domestica* Linnaeus. Folkestone, London, 743 pp.
Included some references applicable to *H. i. irritans*.
393. WIESMANN, R. 1943. Eine neue methode der bekämpfung der fliegenplagen in stallen. Anz. Schaedlingskd. 19(1): 5-8.
Discussed fly control with gesarol (DDT).
394. WILLISTON, S. W. 1896. Manual of the families and genera of the North American Diptera. New Haven, p. 143.
395. WINGO, C. W., G. D. THOMAS, G. N. CLARK, and C. E. MORGAN. 1974. Succession and abundance of insects in pasture manure: relationship to face fly survival. Ann. Entomol. Soc. Am. 67(3): 386-390.
396. WOLFF, A. 1975. Hormones and pheromones. Rockefeller Foundation Illustrated 2(3): 5.
397. ZANGHERI, P. 1950. Ditteri de Romagna. Mem. Soc. Entomol. Ital. 29: 89.
Lyperosia (= *Haematobia*) *irritans* was collected in the Romagna area, northern Italy.
398. ZUMPT, F. 1938. I. Vorstudie zu einer monographischen bearbeitung der Stomoxydinae. Taxonomische ergebnisse des studiums einiger sammlungen, besonders der von Bezzi und Enderlein. Z. Angew. Entomol. 25: 337-353.
Revision of the stomoxyine flies was presented.
399. ——— 1967. *Siphona kangwagyei* n. sp., a bloodsucking fly from Uganda (Diptera: Muscidae). Novos Taxa Entomol. 52, 4 pp.
400. ——— 1969. *Haematobia wooffi* n. sp., and the status of the genus *Bdellia* Enderlein (Diptera: Muscidae, Stomoxyinae). Novos Taxa Entomol. 65, 11 pp.
401. ——— 1973. The Stomoxyine biting flies of the world. Stuttgart, 175 pp.
Presented a revision of the stomoxyine flies of the world. Included morphology and anatomy, classification of various genera and species, life history, and control measures. An extensive list of references was included.
402. ZWALUWENBURG, R. H. VAN. 1928. The interrelationships of insects and roundworms. Hawaii Sugar Plant. Assoc. Exp. Stn. Bull., Entomol. Ser. No. 20, p. 51.



U. S. DEPARTMENT OF AGRICULTURE
AGRICULTURAL RESEARCH SERVICE
HYATTSVILLE, MARYLAND 20782

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, \$300

POSTAGE AND FEES PAID
U. S. DEPARTMENT OF
AGRICULTURE
AGR 101

